

# Immunization Schedule

*Birth:* Hepatitis B

2 Months:

- Pediarix (DTap, hepatitis B, polio)
- Pedvax HIB
- Vaxneuvance 15 (pneumococcal)
- Rotarix (oral vaccination against rotavirus)

4 Months:

- Pediarix (DTap, hepatitis B, polio)
- Pedvax HIB
- Vaxneuvance 15 (pneumococcal)
- Rotarix (oral vaccination against rotavirus)

6 Months

- Pediarix (DTap, hepatitis B, polio)
- Vaxneuvance 15 (pneumococcal)

12 Months:

- MMR (measles, mumps, rubella)
- Varicella (chickenpox)
- Hepatitis A

15 Months:

- Vaxneuvance 15 (pneumococcal)
- Pedvax HIB

18 Months:

- Hepatitis A
- DTap

24 Months:

- No scheduled vaccines/ catch up

4 Years:

- Proquad (MMR, varicella)
- Dtap/IPV

11 Years:

- MenquadFi (meningococcal ACYW)
- Tdap
- Gardasil 9 (HPV)\*

16 Years:

- MenquadFi (meningococcal ACYW)
- Bexsero (meningococcal B)\*

## Highly Recommended Vaccines

\*-**Gardasil 9 (HPV):** Adolescents and teens starting the series before 15 years old only need 2 doses separated by 6 months. Otherwise, it is a 3 dose series.

\*-**Bexsero:** Bexsero is a 2 dose vaccine separated by 30 days. Please be sure to schedule an appointment for dose 2.

## Flu Vaccine

Starting at 6 months of age children should begin receiving an annual flu vaccine. If they are receiving a flu vaccine for the first time, they will need 2 doses separated by one month

---

Please refer to the provided Vaccine Information Statement (VIS) additional information regarding specific vaccines and vaccine preventable illnesses. VIS statements can also be found on the CDC website: [www.cdc.gov](http://www.cdc.gov)